

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 9/127, 31/665, 31/675, 31/685, C07D 259/00, 487/22, C07F 9/02	A1	(11) International Publication Number: WO 00/48572 (43) International Publication Date: 24 August 2000 (24.08.00)
(21) International Application Number: PCT/US00/04140 (22) International Filing Date: 16 February 2000 (16.02.00) (30) Priority Data: 60/120,483 18 February 1999 (18.02.99) US (71) Applicant (for all designated States except US): AMUR PHARMACEUTICALS, INC. [US/US]; 227 Lyndhurst Avenue, Belmont, CA 94002 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MORIMOTO, Bruce, H. [US/US]; 2025 Helena Way, Redwood City, CA 94061 (US). BARKER, Peter, L. [US/US]; 510 Plaza Alhambra #7, El Granada, CA 94018 (US). (74) Agents: FRANKFORT, Howard, M. et al.; Darby & Darby P.C., 805 Third Avenue, New York, NY 10022-7513 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PHOSPHOCHOLINE LINKED PRODRUG DERIVATIVES <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div> (57) Abstract <p>Disclosed are compounds of general formula (I) that function as prodrugs, thereby increasing bioavailabilities of the linked therapeutic agents, wherein the LINKER is (i) substituted or unsubstituted alkyl, (ii) substituted or unsubstituted alkenyl, (iii) substituted or unsubstituted alkanoyl, (iv) substituted or unsubstituted alkenoyl wherein the double bond is <i>cis</i>, and (v) (<i>ortho</i> or <i>para</i>) carbonyl-substituted aryl; and wherein the substituent is each an independent group or linked together thereby forming a ring; and wherein X is one or more substituted or unsubstituted group containing one or more O, N, or S atom and wherein the substituent is each an independent group or linked together thereby forming a ring; and wherein the therapeutic agent is an alcohol-containing water-insoluble steroids or another alcohol containing compounds and methods to prepare such compounds.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	ME	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 00/48572

PCT/US00/04140

PHOSPHOCHOLINE LINKED PRODRUG DERIVATIVESFIELD OF THE INVENTION

The present invention is directed to a novel class of phosphocholine-linked derivatives which not only increase water solubility, but also function as true prodrugs, allow phosphocholines or phosphocholine congeners to be attached to a variety of functional groups on the therapeutic agent, and have the potential on being able to control the rate of release of the pharmaceutical agent.

BACKGROUND OF THE INVENTION

Conventional means for delivering pharmaceutical and therapeutic agents to mammals often are severely limited by chemical and physical properties of the agent, such as aqueous solubility. For example, oral delivery of many biologically-active agents would be the route of choice if not for poor bioavailability due to the limited dissolution of the active agent and subsequent absorption.

Water insoluble therapeutic agents are particularly difficult to administer parenterally. Formulations often require inclusion of a variety of emulsifiers, such as CREMOPHOR® EL. But CREMOPHOR®, which is poly(oxyethylene)-40-castor oil, can result in hypotension, dyspnea, angioedema, or generalized urticaria. These hypersensitive reactions can lead to life-threatening conditions, and it is recommended that all patients be premedicated with corticosteroids, diphenhydramine, and H2 antagonists to avoid severe hypersensitivity.

WO 00/48572

PCT/US00/04140

-2-

Another emulsifier is administered in Propofol, an anesthetic. This emulsion contains soy bean oil, glycerol, and egg phosphatide that create a microbial contamination problem with the current formulation of propofol, which can result in life-threatening illness or death from fever, infection or sepsis. This is especially problematic for post-operative or intensive care unit (ICU) patients. Although U.S. Patent 5,714,120 discloses a method to minimize microbial contamination by the addition of a preservative, this formulation is not an antimicrobial preserved product by USP standards and extrinsic contamination remains problematic.

There is thus a need in the art for methods and compositions to enable potential therapeutic agents to be rendered soluble thereby circumventing the need for emulsifiers and providing for safer and more efficacious therapeutic agents.

SUMMARY OF THE INVENTION:

In one aspect, the present invention provides a method for enabling potential therapeutic agents to be rendered soluble comprising the steps of inserting one or more linker moieties having one or more primary alcohol group between a phosphocholine or a phosphocholine congener to the therapeutic agents having one or more compatible group.

In another aspect, the present invention provides a method for increasing the bioavailability of a pharmaceutical agent, comprising the steps of derivatizing the agent with one or more linker moieties, producing an intermediate, recovering and coupling the intermediate with phosphocholine or a phosphocholine-congener to the linkers, producing a final derivative and administering the final derivative to a mammal, wherein the agent in derivative form is significantly more soluble in aqueous media than the agent in non-derivatized form.

WG 00/48572

PCT/US00/04140

-3-

In yet another aspect, the present invention provides a composition of matter comprising an isolated phosphocholine linked via a linker to propofol, a sedative or anesthetic agent.

5 In yet another aspect, the present invention provides a pharmaceutical formulation for treating a mammal suffering from cancer comprising an isolated phosphocholine linked via a linker to paclitaxel and a physiologically acceptable vehicle, carrier, binder, preservative, stabilizer, flavor, etc., as called for by accepted pharmaceutical practice.

10 The aqueous solubilities of the compounds described herein are evaluated by several methods known in the art, such as preparing a saturated solution of the compound in water, removing a known volume of the solution, and quantitating the amount of the compound in that solution using standard
15 analytical techniques, like HPLC or LC-MS.

These and other aspects of the present invention will be apparent to those of ordinary skill in the art in light of the present description, claims and drawings.

DETAILED DESCRIPTION OF THE INVENTION

20 The invention in its broad aspects relates to phosphocholine or phosphocholine congeners, attached via a linker, to a therapeutic agent.

Phosphocholine derivatives of therapeutic agents containing a primary alcohol or a phenol are readily cleaved by
25 *phosphatases* and mammalian *esterases*. The preparation of phosphocholine derivatives of biologically active agents has been reported (e.g., U.S. Patent No. 5,703,063). However, if the phosphocholine is attached to a secondary or sterically hindered alcohol, hydrolysis or removal of the phosphocholine does not
30 occur rapidly or to a large extent.

WO 00/48572

PCT/US00/04140

-4-

The present invention advantageously provides insertion of a linker between the phosphocholine and the secondary alcohol of the therapeutic agent wherein the phosphocholine is bound to the linker via a primary alcohol or phenol functional group inherent in the linker. This formulation facilitates enzymatic cleavage of the phosphocholine linker bond and liberates the primary alcohol or phenol of the linker. The linker then spontaneously eliminates to liberate the therapeutic agent and an inert molecule arising from the decomposed linker.

Poor water solubility of biologically active agents is, in many cases, the reason for poor bioavailability of the compounds. The compounds described herein display no less than 5 to 10-fold increased biological activity and/or aqueous solubility as compared to the non-derivatized therapeutic agents when administered by the same route. Preferably, the compounds display increased biological activity and/or aqueous solubility in the range from one hundred fold to one hundred thousand fold relative to the non-derivatized therapeutic agents. Thus, the compounds described herein are useful for the enhanced bioavailability of otherwise water-insoluble compounds.

The phosphocholine congeners include, but are not limited to, O-phosphoserine; O-phosphothreonine; O-phosphotyrosine and their mono- and di-N-methyl derivatives; O-phosphoethanolamine and their mono- and di-N-methyl derivatives.


Although the compounds of this invention can include phosphocholine or phosphocholine congeners, they will be described below as compounds having phosphocholine of general FORMULA I:

30

WO 00/48572

PCT/US00/04140

-5-



wherein the LINKER is one or more of the groups selected from the group consisting of (i) substituted or unsubstituted alkyl, (ii) substituted or unsubstituted alkenyl, (iii) substituted or unsubstituted alkanoyl, (iv) substituted or unsubstituted alkenoyl wherein the double bond is *cis*, and (v) (*ortho* or *para*) carbonyl-substituted aryl; and

wherein the substituent is each an independent group or linked together thereby forming a ring; and

wherein X is one or more substituted or unsubstituted group containing one or more O, N, or S atom and

wherein the substituent is each an independent group or linked together thereby forming a ring; and

wherein the therapeutic agent is selected from the group consisting of alcohol-containing water-insoluble steroids and another alcohol containing compounds.

The (*ortho* or *para*) carbonyl-substituted aryl of the LINKER is selected from the group consisting of *ortho*-CR₁R₂-substituted aryl-CO, substituted aryl-*ortho*-CR₃R₄-CO, substituted aryl-*ortho*-CR₃R₄-CR₅R₆-CO, substituted aryl-*ortho*-CR₃=CR₄-CO wherein the double bond is *cis*, *ortho*-CR₁R₂-substituted aryl-CR₅R₆-CO, and substituted aryl-(*ortho* or *para*)-CO.

The aryl substituents may optionally be selected to accelerate or decelerate the rate of enzymatic cleavage of a phenolic phosphocholine. Examples of substituents accelerating the rate of enzymatic cleavage would be nitro, alkyl or aryl sulfonyl, alkyl or aryl keto, alkyl or aryl oxycarbonyl in the

WO 00/48572

PCT/US00/04140

-6-

ortho and/or para positions relative to the phenolic phosphocholine. Examples of substituents decelerating the rate of enzymatic cleavage of a phenolic phosphocholine would be alkyl, alkoxy, alkylthio in the ortho and/or para positions relative to the phenolic phosphocholine.

The aryl is selected from the group consisting of benzene, naphthalene, pyridine, pyrrole, thiophene, furan, imidazole, thiazole, oxazole, pyrimidine, indole, benzimidazole, benzthiazole, benzofuran, benzothiophene and quinoline, each bearing one or more of the group consisting of hydrogen, C₁₋₈-alkyl, C₁₋₈-alkoxy, F, Cl, Br, C₁₋₈-alkoxycarbonyl, amino, substituted amino, nitro, C₁₋₈-alkylthio, C₁₋₈-alkyl sulfoxido, and C₁₋₈-alkylsulfonyl.

In one embodiment, the present invention is a compound having a general formula I wherein (i) said alkyl has the formula CR₁R₂, (ii) said alkenyl has the formula CR₁=CR₃-CR₄, (iii) said alkanoyl has the formula CR₁R₂-CR₃R₄-CR₅R₆-CO, (iv) said alkenoyl has the formula CR₁R₂-CR₃=CR₄-CO and wherein the double bond is *cis*, and (v) said substituted aryl has the formula aryl-CR₁R₂; and

wherein R₁, R₂, R₃, R₄, R₅, and R₆ are the same or different and are selected from the group consisting of

(i) hydrogen;
(ii) linear, branched, and unsaturated C₁₋₁₂-alkyl;
(iii) substituted C₁₋₈-alkyl, wherein the substituent is selected from the group consisting of Y1-Y24, wherein

Y1 is hydroxy,
Y2 is C₁₋₈-alkoxy,
Y3 is carbo-C₁₋₈-alkoxy,
Y4 is C₁₋₈-alkylamino,
Y5 is di-C₁₋₈-alkylamino,
Y6 is C₆₋₁₂-arylamino,
Y7 is C₆₋₁₂-aryloxy,
Y8 is amino,

WO 00/48572

PCT/US00/04140

-7-

- Y9 is amino-C₂-C₈-alkoxy,
Y10 is C₁₋₈-alkylthio,
Y11 is C₆₋₁₂-arylthio,
Y12 is acetamido,
5 Y13 is mercapto,
Y14 is benzamido,
Y15 is carboxamido,
Y16 is phthalimido,
Y17 is guanidino,
10 Y18 is ureido,
Y19 is isothioureido,
Y20 is carboxy,
Y21 is (C₆₋₁₂) aryl-(C₁₋₈) alkyl,
Y22 is (C₆₋₁₂) aryl-(C₂₋₈) alkenyl,
15 Y23 is aromatic heterocyclo(C₁₋₈) alkyl,
and Y24 is aromatic heterocyclo-(C₂₋₈)-alkenyl wherein
the heterocyclic group of Y23 and Y24 have 5-10 ring atoms and
have up to two O, N, or S heteroatoms; and (iv) substituted Y21
or substituted Y23 wherein the substituent is selected from the
20 group consisting of Y1, Y2, Y4, Y5, Y7, Y8, Y12, Y14, Y17-Y20,
and Y25-Y29 wherein
Y25 is halogen,
Y26 is C₁₋₈-alkyl,
Y27 is amino-C₁₋₈-alkyl,
25 Y28 is C₆₋₁₂-aroyl, and
Y29 is C₁₋₈-alkanoyl.

- Unless specified otherwise: (i) alkyl, alkenyl and
alkynyl denote straight and branched hydrocarbon chains having
single, double and triple bonds, respectively; (ii) C₆₋₁₂-aryl
30 groups denote unsubstituted aromatic ring or rings such as, for
example, phenyl or naphthyl; (iii) hetero denotes the
heteroatoms O, N, or S; (iv) aromatic heterocyclic have five to
ten ring atoms and contain up to four heteroatoms; (v) halogen
or halo denote F, Cl, Br, or I atoms; and (vi) alkoxy denotes an
35 alkyl group attached to O.

WO 00/48572

PCT/US00/04140

-8-

Examples of C₁₋₆-alkyl or C₂₋₆ alkenyl groups include methyl, ethyl, propyl, isopropyl, butyl, t-butyl, sec-butyl, pentyl, isopentyl, hexyl, vinyl, allyl, butenyl and the like; aromatic heterocyclic group is selected from the group consisting of pyridyl, thienyl, furyl, indoyl, benzthienyl, imidazolyl, thiazolyl, quinolyl and isoquinoyl.

The compounds containing the R-groups described herein can be purchased from numerous commercial sources such as Sigma Chemical Company (St. Louis, MO), Aldrich Chemical Company (Milwaukee, WI), Acros Organic Chemicals (Pittsburgh, PA), or Fluka Chemical Corporation (Milwaukee, WI). All other compounds not directly available from commercial sources can be prepared from commercially available starting materials by anyone skilled in the art of synthetic organic chemistry.

The preferred linkers are the compounds wherein R₁ is hydrogen, and R₂, R₃, R₄, R₅ and R₆ are the same or different and are selected from the group as defined above.

The most preferred linkers are compounds of the above formula wherein R₁ and R₂ are hydrogen and R₃, R₄, R₅ and R₆ are the same or different and are selected from the group as defined above.

More than one linker per therapeutic agent molecule can be present when more than one appropriate functional group (X) exists. In such case, the order of removal of multiple phosphocholines on a single therapeutic agent would depend on a number of factors: (i) steric effects, (ii) nature of linker, (iii) nature of X. Steric effects influencing the order of removal of multiple phosphocholines would be determined by the immediate steric environment of the specific phosphocholine-linked therapeutic agent. Sterically crowded phosphocholine-linked therapeutic agents would be predicted to be enzymatically cleaved more slowly than non-sterically crowded phosphocholine-linked therapeutic agents. Substituents on the linker may drive the elimination of the linker by sterically favoring a geometric form of the

WO 00/48572

PCT/US00/04140

-9-

intermediate linker-therapeutic agent which self-eliminates more rapidly. Variables in the nature of the linker include inherent differences in the kinetics of the decomposition of the various linkers, and include the nature of substituents of substituted phenyl based linker as described (above/below). Variables in the nature of X include electronic effects of X as a leaving group. Generally, the more electronically deficient X is a better leaving group and hence is eliminated more rapidly and regenerates the therapeutic agent faster than an electronically rich X.

X is selected the group containing one or more O, N, or S atom selected from the group consisting of O, (O)CO, NR₁, NR₂, CO, NR₂ CO NR₃, NR₂(SO₂), NR₂ CS, NR₂ CS NR₃, ONR₂, ONR₂CO, NR₂(O), NR₂(O)CO, nitrogen heterocycles, amide and urea internal in the therapeutic agent.

R₁ and R₂ are the same or different and are selected from the group consisting of

- (i) hydrogen;
- (ii) linear, branched, and unsaturated C₁₋₁₂-alkyl;
- (iii) substituted C₁₋₈-alkyl, wherein the substituent is selected from the group consisting of Y1-Y13 and Y15-Y25;
- (iv) substituted Y21 or substituted Y23 wherein the substituent is selected from the group consisting of Y1, Y2, Y4, Y5, Y7, Y8, Y12, Y14, Y17-Y20, and Y25-Y29.

There may be more than one X in the therapeutic agent and, hence, more than one phosphocholine linked to the therapeutic agent.

R₁ and R₂ may be linked together thereby forming

- (i) a ring of three to six carbon atoms, or
- (ii) a ring of two to five carbon atoms and one O, or S heteroatom, or substituted heteroatom NR₁; wherein R₁ is selected from the group consisting of Y21, Y26, and Y28-Y31.

R₁ and / or R₂ may be connected to the therapeutic agent molecule thereby forming alkylene bridge of from one to five carbon atoms and one or two O, S or NR₁ heteroatoms; wherein R₁ is selected from the group consisting of Y21, Y26,

WO 00/48572

PCT/US00/04140

-10-

Y28-Y31, and the pharmaceutically acceptable salts thereof.

Examples of therapeutic agents which benefit from a phosphocholine linker:

5 Numerous biologically active compounds suffer from low water solubility and poor bioavailability. One family of such compounds are steroids, which are in general, poorly bioavailable. The steroids include testosterone, cardiotonic steroids, and other steroids with biological activity.

10 Testosterone is prescribed therapeutically for men with low levels of endogenous testosterone. Delivery is problematic, however, and necessitates the use of, for example, a testosterone impregnated patch which must be applied directly to the shaven scrotum. A water soluble phosphocholine-linked prodrugs of testosterone could therefore
15 be useful in circumventing delivery of the therapeutic agent.

Cardiotonic steroids, such as digitoxigenin, digoxigenin and ouabugenin are currently used therapeutically. However, their low levels of oral availability makes dosing difficult and the potential for an overdose an important
20 consideration for the attending physician. Phosphocholine linked steroids have the potential to be delivered intravenously, nasally, perorally, intratracheally, administered by patch, etc.

Other steroids with biological activity are
25 candidates for derivatization with phosphocholine linkers. Dehydroepiandrosterone (DHEA), etiocholanolone, pregnenolone, estradiol, estrone, dexamethasone and hydrocortisone are a few examples of steroids which could benefit by deriavatization with a phosphocholine linker.

30 Anti-neoplastic agents, for example, paclitaxel and other taxanes, etoposide, vincristine, vinblastine, and topoisomerase I inhibitors like camptothecin, irinotecan (Pharmacia & Upjohn, Kalamazoo, MI), topotecan (SmithKline Beecham, Philadelphia, PA), CPT11 (Bristol-Myers Squibb,
35 Princeton, NJ); antiviral agents, including nucleoside analogs

WO 00/48572

PCT/US00/04140

-11-

and protease inhibitors, such as nelfinavir (Agouron, LaJolla, CA), saquinavir (Roche, Nutley, NJ), crixivan (Merck, West Point, PA), ritonavir (Abbott, N. Chicago, IL); antibiotics, particularly mitomycin, bleomycin, daunorubicin, doxorubicin, actinomycin, and amphotericin; anesthetics, such as propofol and barbituates, for use in general anesthesia or sedation; analgesics, such as morphine, codeine, and Ziconotide (Neurex, Menlo Park, CA); therapeutic peptides or peptidomimetics, composed of D-amino acids, L-amino acids, or amino acid analogs, acting as enzyme inhibitors, receptor ligands, or disruptors of protein-protein interactions, such as cyclosporin A; therapeutic polypeptides or proteins, such as leptin, growth hormone, calcitonin, vasopressin, renin, prolactin, thyroid and parathyroid hormones, corticotropin, corticotropin-releasing factor, follicle stimulating hormone, luteinizing hormone, gonadotropin, atrial peptides, isolated from natural sources or produced by recombinant DNA technology; nucleic acids, such as anti-sense oligonucleotides or nucleic acids for gene therapy, composed of ribo- or deoxyribonucleotides or nucleotide analogs. Unless otherwise noted, the compounds described herein can be purchased from numerous commercial sources, such as Sigma Chemical Company (St. Louis, MO, Calbiochem-Novabiochem (San Diego, CA), Research Biochemicals Inc. (Natick, MA), or Alexis Corp. (San Diego, CA).

Particularly preferred therapeutic agents for use in the present invention are Propofol and related anesthetic/sedative compounds. These compounds can be conjugated to phosphocholine or phosphocholine congeners via one or more linker pursuant to the present invention and used as anesthetic compounds. It is expected that these derivatized agents will be more effective due to their increased aqueous solubility.

WO 00/48572

PCT/US00/04140

-12-

Compounds of Formula I in pharmaceutical compositions

The derivatized prodrugs of the present invention can be incorporated into pharmaceutical formulations to be used to treat mammals. Pharmaceutical formulations comprising the phosphocholine linked prodrugs derivatives of the present invention as one or more of the active ingredients, would in addition optionally comprise pharmaceutically-acceptable carriers, diluents, fillers, salts and other materials well-known in the art depending upon the dosage form utilized. The compounds of this invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration; suppositories for rectal administration; sterile solutions or suspensions for injectable administration; sterile solutions for ocular or intranasal administration, and the like.

Animals in need of treatment using compounds of this invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from animal to animal and be dependent on such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize.

Typical formulation of compounds of Formula I as pharmaceutical compositions are discussed below. It will be appreciated that the unit content of active ingredient or ingredients contained in an individual dose or dosage form need not in itself constitute an effective amount for the various usages of the phosphocholine linked prodrugs derivatives of the present invention since the necessary effective amount can be reached by administration of a plurality of such dosage forms.

About 0.5 to 100 mg of a compound or mixture of compounds, as the zwitterionic phosphocholine or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, binder, preservative, stabilizer, flavor, etc., as called for by accepted pharmaceutical practice. The amount of active

WO 00/48572

PCT/US00/04140

-13-

ingredients in these compositions is such that a suitable dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules and the like are a binder such as acacia, corn starch or gelatin; an excipient such as microcrystalline cellulose; a disintegrating agent like corn starch or alginic acid; a lubricant such as magnesium stearate; a sweetening agent such as peppermint, wintergreen or cherry. When the dosage form is in a capsule, in addition to the above materials it may also contain a liquid carrier such as a fatty oil.

Other materials of various types may be used as coatings or as modifiers of the physical form of the dosage unit. A syrup or elixir may contain the active compound, a sweetener such as sucrose, preservatives such as propyl paraben, a coloring agent and a flavoring agent such as cherry. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example, dissolution or suspension of the active compound in a vehicle such as water or naturally occurring vegetable oil like sesame, peanut, or cottonseed oil or a synthetic fatty vehicle like ethyl oleate or the like may be desired.

Buffers, preservatives, antioxidants and the like can be incorporated according to the acceptable pharmaceutical practice.

The products of Formula I can be made by using the following general synthetic scheme. The definitions of the substituent groups are the same as for Formula I except where noted. The following examples of reagents are intended to further illustrate the present invention without limiting it thereof.

WG 00/48572

PCT/US00/04140

-14-

GENERAL SYNTHETIC SCHEME

Where Therapeutic agent = alcohol containing water insoluble steroids or another alcohol containing compound as defined in the specification; and


where X = O, N, or S containing groups as defined in the specification; and

where LINKER = unsubstituted or substituted alkyls or phenyls as defined in the specification; and where the primary alcohol part of the LINKER is protected with a tetrahydropyran (THP) or another alcohol protecting groups.


DCC
DMAP
CHCl₃




Dowex (H⁺)



N(CH₂CH₃)₃



N(CH₃)₃



(FORMULA I)


WO 00/48572

PCT/US00/04140

-15-

The preferred products of Formula I can also be made by using the method depicted below. The definitions of the substituent groups are the same as for Formula I except where noted.

5



10

The following examples of compounds of Formula I by using Propofol as a therapeutic agent are illustrated in Example 1 below in Methods A or B. The biological activity of

WG 00/48572

PCT/US00/04140

-16-

phosphocholine linked prodrugs derivatives of the present invention are illustrated in the Example 2. Both examples are intended to further illustrate the present invention without limiting it thereof. The definitions of the substituent groups are the same as for Formula I except where noted.

Example 1

Method A

Preparation of Phosphocholine-linked Propofol (sedative/anesthetic) {2',6'-Diisopropylphenyl 4-(2-trimethyl ammonium ethyloxy)phosphonobutyrate}.

Ethyl 4-hydroxycrotonate (trans) (Kende, Org. Syn. Col. Vol. VII, p221) was treated with 2,3-dihydropyran and catalytic toluenesulfonic acid, according to Bernady (J. Org. Chem. 44, 1438, 1979) to yield ethyl 4-[2-tetrahydro pyranyl]oxycrotonate (trans). This compound was further treated with 0.1M LiOH in tetrahydrofuran, to yield the free acid (4-[2-tetrahydropyranyl] oxycrotonic acid), after acidification and work-up. This carboxylic acid was then coupled, via an ester bond, to 2,6-diisopropylphenol, utilizing *N,N*-dicyclohexyl-carbodiimide. After chromatographic purification on silica gel, the ester was then treated, in methanol, with a catalytic amount of Dowex 50W ion exchange resin to affect the removal of the tetrahydropyranyl protecting group. The resulting alcohol was treated with 2-chloro-2-oxo-1,3,2-dioxaphospholane in the presence of triethylamine in chloroform. Upon completion of this reaction, the isolated phosphate intermediate was dissolved in acetonitrile, charged with trimethylamine, and heated @ 80°C for 72 hours. After removal of trimethylamine, the solvent was removed in vacuo, the residue was partitioned between ethyl acetate and water. Freezing and lyophilization of the aqueous phase yielded the 4-O-phosphocholine. This compound was then hydrogenated in water to yield the title compound. LC/MS, NMR, and combustion data are available.

WO 00/48572

PCT/US00/04140

-17-

Method B

Preparation of Phosphocholine-linked Propofol (Sedative, anesthetic) {2',6'-Diisopropylphenyl 3-[ortho-(O-trimethylammoniummethyl phosphonooxy)]propionate}.

5 To a solution of 3g of 2-hydroxycinnamic acid (trans) in 60mL of dry chloroform was added 7.6mL (eq) of triethylamine. This solution was cooled in an ice bath and 5.7g (2.2eq) of 2-chloro-1,3,5-dioxaphospholane-2-oxide was added dropwise at 0°C. The reaction was allowed to stir at
10 room temperature for thirty minutes. A solution of 2,6 diisopropylphenol (Propofol) in 20mL of chloroform was added and the reaction was stirred at room temperature for 16hr. The reaction was then washed three times with water, and then dried (MgSO₄). Filtration and evaporation of the solvent
15 yielded 10.5g of crude 2',6'-diisopropylphenyl 3-[ortho-(O-ethylene phosphonooxy)]propionate. This intermediate was treated with excess trimethylamine in acetonitrile in a pressure vessel at 80°C for 72hr. Removal of the excess trimethylamine and evaporation of the acetonitrile yielded the
20 crude phosphocholine derivative of 2',6'- diisopropyl phenyl 2-hydroxycinnamate. This intermediate was purified by chromatography (silica gel, CHCl₃/MeOH/H₂O 40:55:5) yielding approximately 100mg of material. Hydrogenation of this intermediate in aqueous ethanol, employing 5%Pd/C yielded the
25 title compound. LC/MS and NMR data are available.

Example 2Sleep indication in mice

The method which detects sedative activity following the protocol described by Simon et al. (*J. Pharmacol. Paris*,
30 13:241-252, 1982).

Mice (10 per group) are placed in Plexiglass cages (20 x 10 x 10 cm) and administered the test substance, propofol, produced as above as an i.v. bolus in two seconds. The latency to sleep and the occurrence of sedation/sleep are
35 noted over a period of one hour. Sleep is indicated by the

WO 00/48572

PCT/US00/04140

-18-

loss of the righting reflex. Animals within a group are tested sequentially and the test is performed blind. The test substance will be evaluated in 5 escalating doses. Unmodified propofol (16 mg per kg) administered in the same experimental conditions, will be used as a reference compound. The LD₅₀ for hypnotic activity is calculated following the method of Lichtfield and Wilcoxin (*J. Pharmacol. Exp. Ther.* 96:99-113, 1949).

Lethal dose 50 (LD₅₀) in mice

The method, which determines the acute dose of a test substance causing 50% of death in a given animal species, follows the method described by Lichtfield and Wilcoxin.

After an 18 hour period of food deprivation but free access to water, mice will be administered the test substance as an i.v. bolus in two seconds. The appearance of morbidity, including local reaction and mortality, are noted for a period of 7 days, during which the animals have free access to food and water.

Ten mice are studied per group. The test is performed blind.

The test substance will be evaluated at 5 escalating doses.

No reference substance and no control group are offered for this experiment.

The LD₅₀ is calculated at the end of the testing following the method of Lichtfield and Wilcoxin see *J. Pharmacol. Exp. Ther.* 96:99-113, 1949 above.

Variations of the present invention will suggest themselves to those skilled in the art, and are within the scope of the following claims:


WO 00/48572

PCT/US00/04140

-19-

We Claim:

- 1 1. A compound having the general formula I:



- 2 wherein the LINKER is one or more of the groups
3 selected from the group consisting of (i) substituted or
4 unsubstituted alkyl, (ii) substituted or unsubstituted alkenyl,
5 (iii) substituted or unsubstituted alkanoyl, (iv) substituted or
6 unsubstituted alkenoyl wherein the double bond is *cis*, and
7 (v) (*ortho* or *para*) carbonyl-substituted aryl; and
8 wherein the substituent is each an independent group or
9 linked together thereby forming a ring; and
10 wherein X is one or more substituted or unsubstituted
11 group containing one or more O, N, or S atom and
12 wherein the substituent is each an independent group or
13 linked together thereby forming a ring; and
14 wherein the therapeutic agent is selected from the
15 group consisting of alcohol- containing water-insoluble steroids
16 and another alcohol containing compounds.

- 1 2. A compound according to claim 1, wherein
2 (i) said alkyl has the formula CR₁R₂, (ii) said alkenyl has the
3 formula CR₁=CR₃-CR₄, (iii) said alkanoyl has the formula CR₁R₂-
4 CR₃R₄-CR₅R₆-CO, (iv) said alkenoyl has the formula CR₁R₂-CR₃=CR₄-CO
5 and wherein the double bond is *cis*, and (v) said substituted aryl
6 has the formula aryl-CR₁R₂; and
7 wherein R₁, R₂, R₃, R₄, R₅, and R₆ are the same or

WO 00/48572

PCT/US00/04140

-20-

8 different and are selected from the group consisting of
9 (i) hydrogen;
10 (ii) linear, branched, and unsaturated C₁₋₁₂-alkyl;
11 (iii) substituted C₁₋₈-alkyl, wherein the substituent is
12 selected from the group consisting of Y1-Y24, wherein
13 Y1 is hydroxy,
14 Y2 is C₁₋₈-alkoxy,
15 Y3 is carbo-C₁₋₈-alkoxy,
16 Y4 is C₁₋₈-alkylamino,
17 Y5 is di-C₁₋₈-alkylamino,
18 Y6 is C₆₋₁₂-arylamino,
19 Y7 is C₆₋₁₂-aryloxy,
20 Y8 is amino,
21 Y9 is amino-C₂-C₈-alkoxy,
22 Y10 is C₁₋₈-alkylthio,
23 Y11 is C₆₋₁₂-arylthio,
24 Y12 is acetamido,
25 Y13 is mercapto,
26 Y14 is benzamido,
27 Y15 is carboxamido,
28 Y16 is phthalimido,
29 Y17 is guanidino,
30 Y18 is ureido,
31 Y19 is isothioureido,
32 Y20 is carboxy,
33 Y21 is (C₆₋₁₂)aryl-(C₁₋₈)alkyl,
34 Y22 is (C₆₋₁₂)aryl-(C₂₋₈)alkenyl,
35 Y23 is aromatic heterocyclo(C₁₋₈)alkyl,
36 and Y24 is aromatic heterocyclo(C₂₋₈)alkenyl wherein
37 the heterocyclic group of Y23 and Y24 have 5 - 10 ring atoms and
38 comprises up to two O, N, or S heteroatoms; and
39 (iv) substituted Y21 or substituted Y23 wherein the
40 substituent is selected from the group consisting of
41 Y1, Y2, Y4, Y5, Y7, Y8, Y12, Y14, Y17-Y20, and Y25-Y29 wherein

WO 00/48572

PCT/US00/04140

-21-

42 Y25 is halogen,
43 Y26 is C₁₋₈-alkyl,
44 Y27 is amino-C₁₋₈-alkyl,
45 Y28 is C₆₋₁₂-aroyl, and
46 Y29 is C₁₋₈-alkanoyl.

1 3. A compound according to claim 2, wherein said
2 R₁ and R₂; R₁ and R₃; R₂ and R₃; R₃ and R₄; R₃ and R₅; and R₅ and R₆
3 are linked together thereby forming:

4 (i) a ring of three to six carbon atoms, or
5 (ii) a ring of two to five carbon atoms and one O, or S
6 heteroatom, or substituted heteroatom NR₇, wherein R₇ is selected
7 from the group consisting of Y21, Y26, Y28, Y29, and Y30-Y31,
8 wherein Y30 is C₃₋₈-alkenyl, and
9 Y31 is C₆₋₁₂-aryl.

1 4. A compound according to claim 2 wherein the group
2 containing one or more O, N, or S atom is selected from the group
3 consisting of O, (O)CO, NR₈, NR₈CO, NR₈CO NR₉, NR₈(SO₂), NR₈CS,
4 NR₈CS NR₉, ONR₈, ONR₈CO, NR₈(O), NR₈(O)CO, nitrogen heterocycles,
5 amide and urea internal in therapeutic agent; and
6 wherein R₈ and R₉ are the same or different and are
7 selected from the group consisting of
8 (i) hydrogen;
9 (ii) linear, branched, and unsaturated C₁₋₁₂-alkyl;
10 (iii) substituted C₁₋₈-alkyl, wherein the substituent is
11 selected from the group consisting of Y1-Y13 and Y15-Y25;
12 (iv) substituted Y21 or substituted Y23 wherein the
13 substituent is selected from the group consisting of Y1, Y2, Y4,
14 Y5, Y7, Y8, Y12, Y14, Y17-Y20, and Y25-Y29.

1 5. A compound according to claim 4 wherein R₈ and R₉
2 are linked together thereby forming
3 (i) a ring of three to six carbon atoms, or

WO 00/48572

PCT/US00/04140

-22-

4 (ii) a ring of two to five carbon atoms and one O, or S
5 heteroatom, or substituted heteroatom NR₇; wherein R₇ is selected
6 from the group consisting of Y21, Y26, and Y28-Y31.

1 6. A compound according to claim 4 wherein R₈, R₉, or
2 both are connected to the therapeutic agent molecule thereby
3 forming alkylene bridge of from one to five carbon atoms and
4 one or two O, S or NR, heteroatoms; wherein R₇ is selected from
5 the group consisting of Y21, Y26, Y28-Y31, and the
6 pharmaceutically acceptable salts thereof.

1 7. A compound according to claim 5 wherein R₈, R₉, or
2 both are connected to the therapeutic agent molecule thereby
3 forming alkylene bridge of from one to five carbon atoms and
4 one or two O, S or NR, heteroatoms; wherein R₇ is selected from
5 the group consisting of Y21, Y26, Y28-Y31; and the
6 pharmaceutically acceptable salts thereof.

1 8. A compound according to claim 2, wherein said
2 (ortho or para) carbonyl-substituted aryl is selected from the
3 group consisting of ortho-CR₁R₂-substituted aryl-CO, substituted
4 aryl-ortho-CR₃R₄-CO, substituted aryl-ortho-CR₃R₄-CR₅R₆-CO,
5 substituted aryl-ortho-CR₃=CR₄-CO wherein the double bond is *cis*,
6 ortho-CR₁R₂-substituted aryl-CR₅R₆-CO, and substituted aryl-(ortho
7 or para)-CO.

1 9. A compound according to claim 2, wherein said aryl
2 is selected from the group consisting of benzene, naphthalene,
3 pyridine, pyrrole, thiophene, furan, imidazole, thiazole,
4 oxazole, pyrimidine, indole, benzimidazole, benzthiazole,
5 benzofuran, benzothiophene and quinoline, each bearing one or
6 more of the group consisting of hydrogen, C₁₋₈-alkyl, C₁₋₈-alkoxy,
7 F, Cl, Br, C₁₋₈-alkoxycarbonyl, amino, substituted amino, nitro,

WO 00/48572

PCT/US00/04140

-23-

8 C₁₋₈-alkylthio, C₁₋₈-alkylsulfoxido, and C₁₋₈-alkylsulfono.

1 10. A compound according to claim 2, wherein R₁ is
2 hydrogen.

1 11. A compound according to claim 2, wherein R₁ and
2 R₂ are hydrogen.

1 12. A compound according to claim 1, wherein the
2 therapeutic agent is selected from the group consisting of
3 Propofol and related anesthetic or sedative compounds.

1 13. A compound according to claim 1, wherein said
2 water-insoluble steroids are selected from the group consisting
3 of (i) testosterone, (ii) cardiotonic steroids selected from the
4 group consisting of digitoxigenin, digoxigenin and
5 ouabugenin, (iii) dehydroepiandrosterone (DHEA), (iv)
6 etiocholanolone, (v) pregnenolone, (vi) estradiol, (vii) estrone,
7 (viii) dexamethasone and (ix) hydrocortisone.

1 14. A compound according to claim 1, further comprises
2 one or more of the ingredients selected from the group consisting
3 of pharmaceutically-acceptable carriers, diluents, fillers,
4 salts, buffers, preservatives, antioxidants, a binder, an
5 excipient, a disintegrating agent, a lubricant, and a sweetening
6 agent.


1 15. A compound according to claim 1 incorporated into
2 tablets, capsules or elixirs for oral administration;
3 suppositories for rectal administration; sterile solutions or
4 suspensions for injectable administration; sterile solutions for
5 ocular (?) or intranasal administration.

WO 00/48572

PCT/US00/04140


-24-

1 16. A compound having the general formula I:



4 wherein the LINKER is a substituted alkenyl of formula CR_1R_2-
5 $CR_3=CR_4-CO$, wherein R_1 , R_3 , and R_4 are hydrogen and wherein the
6 double bond is trans, and
7 wherein X is O and
8 wherein the therapeutic agent is 2',6'-diisopropyl
9 phenol.

1 17. A compound having the general formula I:



4 wherein the LINKER is a substituted alkanoyl of formula CR_1R_2-
5 $CR_3R_4-CR_5R_6-CO$, wherein R_1, R_2, R_3, R_4, R_5 , and R_6 are H, and
6 wherein X is O and

WO 00/48572

PCT/US00/04140

-25-

1 wherein the therapeutic agent is 2',6'-diisopropyl
2 phenol.

1 18. A method for enabling potential therapeutic agents
2 to be rendered soluble comprising the steps of inserting one or
3 more linker moieties having one or more primary alcohol group
4 between a phosphocholine or a phosphocholine congener to the
5 therapeutic agents having one or more alcohol group.

1 19. A method for increasing the bioavailability of
2 pharmaceutical agent comprising the steps of derivatizing the
3 agent with one or more linker moieties, producing an
4 intermediate, recovering and coupling the intermediate with
5 phosphocholine or a phosphocholine-congener to the linkers,
6 producing a final derivative and administering the final
7 derivative to a mammal, wherein the agent in derivative form is
8 significantly more soluble in aqueous media than the agent in
9 non-derivatized form.

1 20. The method of claim 19 wherein the pharmaceutical
2 agent is propofol.

1 21. A pharmaceutical formulation for treating a mammal
2 suffering from cancer comprising an isolated phosphocholine
3 linked via a linker to paclitaxel and a physiologically
4 acceptable vehicle, carrier, binder, preservative, stabilizer, or
5 flavor as called for by accepted pharmaceutical practice.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/04140

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 9/127, 31/665, 31/675, 31/685; C07D 259/00, 487/22; C07F 9/02
 US CL : 424/450; 514/77, 79, 80, 81, 82, 85, 86, 92, 99, 100; 540/456, 460, 478; 544/243, 337; 546/23; 548/113,
 119; 549/220, 221, 222; 552/506, 507; 558/170, 171, 174

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 U.S. : Please See Continuation Sheet

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,830,432 A (CHASALOW) 03 November 1998.	1-21
A	WO 98/11906 A1 (AMUR PHARMACEUTICALS, INC.) 26 March 1998.	1-21

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another claim or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"A"

document member of the same patent family

Date of the actual completion of the international search

10 May 2000 (10.05.2000)

Date of mailing of the international search report

09 JUN 2000

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Mukund Shah

Telephone No. 703-308-1235

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/04140

Continuation of B. FIELDS SEARCHED Item 1: 424/450; 514/77, 79, 80, 81, 82, 85, 86, 92, 99, 100; 540/456, 460, 478; 544/243, 337; 546/23; 548/113, 119; 549/220, 221, 222; 552/506, 507; 558/170, 171, 174

Form PCT/ISA/210 (extra sheet) (July 1998)